arXiv:1305.0537v1 [math.AG] 2 May 2013

BIRATIONAL GEOMETRY OF HYPERSURFACES IN PRODUCTS OF PROJECTIVE SPACES

JOHN CHRISTIAN OTTEM

Abstract. We study the birational structure of hypersurfaces in products of projective spaces. These hypersurfaces are in many respects simple varieties, yet they provide many interesting examples of birational geometry phenomena. In the case of hypersurfaces in $\mathbb{P}^m \times \mathbb{P}^n$, we describe their nef, movable and effective cones and determine when they are Mori dream spaces.

1. Introduction

Let X be a hypersurface of $\mathbb{P}^m \times \mathbb{P}^n$ of bidegree (d, e) defined by a bihomogeneous polynomial in the variables $x_0, \dots, x_m, y_0, \dots, y_n$. If the dimension of X is at least three, the Lefschetz theorem says that the inclusion induces an isomorphism of Picard groups $\operatorname{Pic}(X) \simeq \operatorname{Pic}(\mathbb{P}^m \times \mathbb{P}^n) \simeq \mathbb{Z}^2$. It is therefore natural to ask how the various cones of divisors of X are related to those of the ambient space $\mathbb{P}^m \times \mathbb{P}^n$. In general this relation is not clear, as examples of Hassett–Lin–Wang [7] show that the nef cone of X can be strictly greater than that of $\mathbb{P}^m \times \mathbb{P}^n$ in the cases $m = 1$ or $n = 1$ (however, for $m, n \geq 2$, the two nef cones coincide). We therefore ask:

Question 1. What are the cones of effective, movable or nef divisors on X?

Recall that a normal $\mathbb{Q}$-factorial projective variety X is a *Mori dream space* if the following three conditions are fulfilled: (i) $H^1(X, \mathscr{O}_X) = 0$; (ii) the nef cone $\operatorname{Nef}(X)$ is generated by the classes of finitely many semiample divisors; and (iii) there is a finite collection of small $\mathbb{Q}$-factorial modifications $\phi_i : X \dashrightarrow X_i$ such that each X_i satisfies (ii) and the movable cone of X decomposes as $\operatorname{Mov}(X) = \bigcup \phi_i^* \operatorname{Nef}(X_i)$.

This notion was introduced by Hu and Keel in [8] as a class of varieties with good birational geometry properties. It turns out that the condition of being a Mori dream space is equivalent to having a finitely generated Cox ring [8, Theorem 2]. This ring is essentially defined by

$$\mathcal{R}(X) = \bigoplus_{D \in \operatorname{Pic} X} H^0(X, \mathscr{O}_X(D)) \tag{1.1}$$

where the ring structure is given by viewing sections as rational functions.

Mori dream spaces are particularly nice from the viewpoint of Mori theory; the Minimal Model Program can be carried out for any divisor and has a combinatorial structure as in the case of toric varieties. More precisely, choosing a presentation for the Cox ring gives an embedding of X into a simplicial toric variety Y such that $\operatorname{Eff}(Y) = \operatorname{Eff}(X)$ and $\operatorname{Nef}(X) = \operatorname{Nef}(Y)$ and each of the modifications ϕ_i above is induced from a small modification of the ambient space Y (see [8, Proposition 2.11]). It is therefore natural to ask:

Question 2. For which d, e is the hypersurface $X \subset \mathbb{P}^m \times \mathbb{P}^n$ a Mori dream space?

Being a Mori dream space is a relatively strong condition and there are classical examples of varieties that are not. Perhaps the most famous of these is Nagata's counterexample to Hilbert's 14th problem, in which he proves that the blow-up of $\mathbb{P}^2$ along the base-locus of a general cubic pencil has infinitely many (-1)-curves [15]. This blow-up is clearly not a Mori dream space since each of the (-1)-curves would require a generator of the Cox ring. The same phenomenon happens for a K3 surface with large Picard number, where one typically expects infinitely many (-2)-curves.

There are also other obstructions to being a Mori dream space than the non-polyhedrality of the effective/nef cones. Indeed, there might be integral classes in the boundary of the nef cone which are not semiample. In this paper we will construct concrete examples of this phenomenon even for Picard number 2; In fact 'most' hypersurfaces in $\mathbb{P}^1 \times \mathbb{P}^n$ have an extremal ray of the nef cone which does not even have an effective multiple, so they are not Mori dream spaces. Thus these hypersurfaces provide simple counterparts to Nagata's examples above.

Our main result is the following:

Theorem 1.1. *Let X be a smooth hypersurface of bidegree (d,e) in $\mathbb{P}^m \times \mathbb{P}^n$ of dimension at least three and let $H_i = \mathrm{pr}_i^* \mathscr{O}(1)$. If $m,n \geq 2$, X is a Mori dream space and $\mathcal{R}(X)$ is isomorphic to $k[x_0,\dots,x_m,y_0,\dots,y_n]/(f)$. In particular,*

$$\mathrm{Eff}(X) = \mathrm{Mov}(X) = \mathrm{Nef}(X) = \mathbb{R}_{\geq 0}H_1 + \mathbb{R}_{\geq 0}H_2$$

For general hypersurfaces in $\mathbb{P}^1 \times \mathbb{P}^n$ where $n \geq 3$, we consider the the following cases:

(i) $d=1$. *Then*

$$\begin{aligned} \mathrm{Eff}(X) &= \mathbb{R}_{\geq 0}H_1 + \mathbb{R}_{\geq 0}(eH_2 - H_1). \\ \mathrm{Mov}(X) &= \mathrm{Nef}(X) = \mathbb{R}_{\geq 0}H_1 + \mathbb{R}_{\geq 0}H_2. \end{aligned}$$

Moreover, the second projection gives X the structure of a blow-up of $\mathbb{P}^n$ along the intersection of two degree e hypersurfaces the exceptional divisor is linearly equivalent to $eH_2 - H_1$. In particular, X is a Mori dream space.

(ii) If $1 < d < n$. *Then*

$$\begin{aligned} \mathrm{Eff}(X) &= \mathrm{Mov}(X) = \mathbb{R}_{\geq 0}H_1 + \mathbb{R}_{\geq 0}(eH_2 - H_1). \\ \mathrm{Nef}(X) &= \mathbb{R}_{\geq 0}H_1 + \mathbb{R}_{\geq 0}H_2. \end{aligned}$$

Moreover, H_2 determines a small contraction $f : X \to Z$ and there is a flip $f^+ : X^+ \to Z$ which gives a decomposition $\mathrm{Mov}(X) = \mathrm{Nef}(X) \cup \mathrm{Nef}(X^+)$. *Hence X is a Mori dream space.*

(iii) $d=n$. *Then*

$$\mathrm{Eff}(X) = \mathrm{Mov}(X) = \mathrm{Nef}(X) = \mathbb{R}_{\geq 0}H_1 + \mathbb{R}_{\geq 0}(eH_2 - H_1).$$

The extremal divisor $eH_2 - H_1$ is base-point free and defines a contraction to $\mathbb{P}^{n-1}$. In particular, X is a Mori dream space.

(iv) If $d > n$, and X is very general, then X is Mori dream if and only if $e = 1$. If $e=1$, X is a $\mathbb{P}^{n-1}$-bundle over $\mathbb{P}^1$. If $e \geq 2$, we have

$$\overline{\mathrm{Eff}(X)} = \overline{\mathrm{Mov}(X)} = \mathrm{Nef}(X) = \mathbb{R}_{\geq 0}H_1 + \mathbb{R}_{\geq 0}(neH_2 - dH_1).$$

But $\mathrm{Eff}(X)$ *is not closed. Hence X is not a Mori dream space.*

Moreover, in Theorem 3.2 below, we give an explicit presentation of the Cox ring of X when it is finitely generated.

Note that the Mori dream hypersurfaces have bidegrees (d,e) lying in the L-shaped region given by $\{(d,e) \in \mathbb{N}^2 \mid d \leq n \text{ or } e = 1\}$. Hence it is essentially the value of d that determines whether a general hypersurface of degree (d,e) is a Mori dream space or not. Note also that all the cones involved are rational polyhedral for any bidegree.

A general principle of Mori theory is that varieties with negative canonical bundle has a better understood birational structure (e.g., the cone theorem gives information about the

K_X-negative part of the cone of curves). It is therefore interesting to interpret Theorem 1.1 in terms of the canonical bundle of X, which is given by $K_X = (d-m-1)H_1 + (e-n-1)H_2$.

Let us for simplicity consider the first non-trivlal case of the theorem, namely hypersurfaces in $\mathbb{P}^1 \times \mathbb{P}^3$. Here there are only a few values of d, e where X is Fano, i.e., when $-K_X$ is ample (using Theorem 1.1, one can show that this happens precisely when $d \leq 2, e \leq 3$; or $e = 1$). In this case it is well-known that X is a Mori dream space [4]. On the other hand, when $(d, e) = (2, 4)$, X is a Calabi-Yau threefold, where $K_X \simeq \mathscr{O}_X$ and $\kappa(X) = 0$, and X is a Mori dream space by the theorem. On the other hand, for general type varieties, when K_X is ample, most hypersurfaces are not Mori dream, although there are some that are (e.g., $(3, 5)$ is Mori dream, $(4, 5)$ is not).

There is a corresponding result for surfaces in $\mathbb{P}^1 \times \mathbb{P}^2$, but this requires a slightly modified argument, as the Picard group of X might be larger than that of the ambient space. Nevertheless, using the Noether-Lefschetz theorem, we prove the following analogue of Theorem 1.1 for surfaces:

Proposition 1.2. *Let X be a very general surface in $\mathbb{P}^1 \times \mathbb{P}^2$ of bidegree (d, e) with $e \geq 2$. We have the following cases:*

- *(i)* $d = 1$. *X is the blow-up of $\mathbb{P}^2$ along the base-locus of two very general degree e curves. It is a Mori dream space if and only if $e \leq 2$, in which case X is a Del Pezzo surface. In the other case, X is a rational surface with infinitely many (-1)-curves.*
- *(ii)* $d = 2$. *X is a double cover of $\mathbb{P}^2$ branched along a smooth curve of degree $2e$ and is always a Mori dream space.*
- *(iii)* $e = 1$. *$X \simeq \mathbb{F}_0$ for d even, and $X \simeq \mathbb{F}_1$ for d odd.*
- *(iv)* $d \geq 3$ *and* $e \geq 2$. *Then the effective cone of X is not closed. Hence X is not Mori dream.*

After this paper was written, we learned that some of the hypersurfaces in Theorem 1.1 were considered by Ito in [9]. More precisely, Ito shows that the hypersurfaces in the cases (i)-(iii) are Mori dream spaces using a different argument than ours.

1.1. **Relation to an example of Mumford.**

Theorem 1.1 gives a simple example of a line bundle L which has positive intersection with every curve C, but is not ample. More precisely, a very general surface of bidegree $(3, 3)$ in $\mathbb{P}^1 \times \mathbb{P}^2$ has the line bundle $L = \mathscr{O}(2H_2 - H_1)$ which satisfies this condition. Of course here L has self-intersection 0. In Section 5 we also construct higher dimensional analogues of this. Examples of line bundles with such properties were first constructed by Mumford using certain projective bundles $\mathbb{P}(\mathscr{E})$ over curves of genus ≥ 2. Our geometric construction in Theorem 1.1(*iv*) was inspired by Mumford's example, but we use only projective bundles over elliptic curves.

1.2. **Relation to the work of Artebani–Laface, Hausen and Jow.**

Motivated by the Lefschetz hyperplane theorem for $\mathrm{Pic}(X)$, a natural question is whether a sufficiently general hypersurface in a Mori dream space is again a Mori dream space. Of course, Theorem 1.1 provides many counterexamples to this, but still it is worthwhile to mention some positive results in this direction.

Let $i : X \hookrightarrow Y$ be an ample hypersurface in a projective Mori dream space Y. Denote by $\overline{Y} = \mathrm{Spec}\, \mathcal{R}(Y)$, and $\overline{X} = V(f) \subset \overline{Y}$ where f is a section defining X in Y. By building on work of Hausen, Artebani–Laface and Jow give criteria for when the restriction of sections induces an isomorphism of Cox rings $i^* : \mathcal{R}(X) \simeq \mathcal{R}(Y)/(f)$. In [10], Jow proves that this holds for any smooth ample divisor X on Y, provided Y is smooth of dimension ≥ 4, and $V(I_{irr}(Y))$ has codimension at least 3 in $\overline{Y}$. Here I_{irr} denotes the so-called *irrelevant ideal*, defined as the radical of the ideal generated by the subalgebra $\bigoplus_{m>0} H^0(Y, mD)$ of $\mathcal{R}(Y)$, where D is any ample divisor on Y. The importance of this ideal comes from the fact that Y is a GIT quotient of $\overline{Y}$ by the Picard torus $\mathrm{Hom}(\mathrm{Pic}(Y), k^*) \simeq \mathbb{G}_m^\rho$ and $V(I_{irr})$ defines the unstable locus of this action [8].

Generalising Jow's results, Artebani and Laface prove very refined results that in particular imply that the above conclusion holds under much weaker assumptions on Y provided X is ample and general in its linear system so that $i^* : \mathrm{Cl}(Y) \simeq \mathrm{Cl}(X)$.

Let us see how these results apply to our situation, where $Y = \mathbb{P}^m \times \mathbb{P}^n$. Here the Cox ring is given by $\mathcal{R}(Y) = k[x_0, \dots, x_m, y_0, \dots, y_n]$, where the grading is $\deg x_i = H_1$ and $\deg y_j = H_2$ and $I_{irr} = (x_0, \dots, x_m) \cap (y_0 \dots, y_n)$. Then it is clear that $V(I_{irr})$ has codimension ≥ 3 if and only if $m, n \geq 2$. Hence when $m, n \geq 2$, Questions 1 and 2 have a straightforward answer:

Proposition 1.3. *Let $m, n \geq 2$ and let $X \subset Y = \mathbb{P}^m \times \mathbb{P}^n$ be a smooth ample hypersurface defined by a general form f of bidegree and let $H_i = \mathrm{pr}_i^* \mathscr{O}(1)$. Then*

$$\mathcal{R}(X) \simeq \mathcal{R}(Y)/(f) = k[x_0, \dots, x_m, y_0, \dots, y_n]/(f)$$

where the bigrading is given by $\deg x_i = H_1$ and $\deg y_i = H_2$. In particular,

$$\mathrm{Nef}(X) = \mathrm{Eff}(X) = \mathrm{Mov}(X) = \mathbb{R}_{\geq 0} H_1 + \mathbb{R}_{\geq 0} H_2$$

Here the same proof works for hypersurfaces in any product of projective spaces of dimension at least 2 (see Corollary 2 of [10]).

In particular, means that the questions mentioned in the introduction are only interesting for hypersurfaces in $\mathbb{P}^1 \times \mathbb{P}^n$. Here the situation becomes more complicated, because the presence of higher cohomology makes it difficult to compute $H^0(X, \mathscr{O}(D))$. In particular, we will see that the conclusion of Proposition 1.3 does not hold in these cases.

1.3. **Notation.** Throughout the paper we will be working over an algebraically closed field of characteristic 0. The main reason for this is that we will use properties of semistable vector bundles (such as the fact that a symmetric power of a semistable vector bundle is again semistable), which are known to be false in positive characteristic. On the other hand it is likely that many of the results in Section 2 and 3 can be extended to positive characteristic using the Grothendick-Lefschetz theorem, which is known to hold in all characteristics.

We let $N^1(X)$ denote the Néron-Severi group of X, i.e., the $\mathbb{R}$-vector space of divisors modulo numerical equivalence. For most of the varieties in this paper numerical and linear equivalence coincide, so that $N^1(X) = \mathrm{Pic}(X) \otimes \mathbb{R}$. Inside $N^1(X)$ we define the effective cone $\mathrm{Eff}(X)$ to be the cone spanned of all effective divisors. Similarly, we denote by $\mathrm{Nef}(X)$ (resp. $\mathrm{Mov}(X)$) the cone of nef divisors (resp. movable divisors). Here we call a divisor D *nef* (resp. *movable*) if $D \cdot C \geq 0$ for every curve C (resp. if the linear system $|D|$ has no fixed components). Note that the nef cone is always closed, whereas the other two need not be.

We will follow the convention of [8] for flips : If $f : X \to Z$ is a small contraction and D is a $\mathbb{Q}$-Cartier divisor such that $-D$ is f-ample, then the *D-flip* of f is a contraction $f^+ : X^+ \to Z$ such that the strict transform of D is $\mathbb{Q}$-Cartier and f^+ ample.

For projectivized bundles, we will use the Grothendieck notation for $\mathbb{P}(\mathscr{E})$, i.e., this is the variety of hyperplanes in the vector bundle $\mathscr{E}$.

We will also distinguish between *general* and *very general* hypersurfaces. Here the former will refer to the hypersurface being chosen outside a finite union of closed algebraic subsets of the parameter space, whereas 'very general' means outside a countable union of closed algebraic subsets.

1.4. **Acknowledgements.** Thanks to my advisor Burt Totaro for his advice and encouragement. Also thanks to Laurent Gruson, Antonio Laface, Victor Lozovanu, Christian Peskine and Claire Voisin for interesting discussions and comments. Special thanks to Kenji Oguiso for sending me his note about threefolds in $\mathbb{P}^1 \times \mathbb{P}^3$.

2. Mori dream hypersurfaces in $\mathbb{P}^1\times\mathbb{P}^n$

Consider a hypersurface $X\subset\mathbb{P}^1\times\mathbb{P}^n$ defined by a bihomogeneous form

$$f=x_0^df_0+x_0^{d-1}x_1f_1+\ldots+x_1^df_d=0 \tag{2.1}$$

where x_0,x_1 coordinates on $\mathbb{P}^1$ and the f_i are homogenous forms of degree e in the coordinates $y_0,\ldots,y_n$ on $\mathbb{P}^n$. We will in this section assume that f is general and that $n\geq 3$. For the case where X is a surface, see Section 4.

Note that X comes with two projections, $\mathrm{pr}_1:X\to\mathbb{P}^1$ and $\mathrm{pr}_2:X\to\mathbb{P}^n$. By the Grothendieck-Lefschetz theorem, we have $\mathrm{Pic}(X)\simeq\mathrm{Pic}(\mathbb{P}^1\times\mathbb{P}^n)=\mathbb{Z}H_1\oplus\mathbb{Z}H_2$ where $H_i=\mathrm{pr}_i^*L_i$ for general hyperplanes $L_1\subset\mathbb{P}^1$ and $L_2\subset\mathbb{P}^n$. We may assume that H_1,H_2 are smooth prime divisors.

We first consider hypersurfaces X of bidegree $(d,1)$ in $\mathbb{P}^1\times\mathbb{P}^n$ with $d\leq n$. After a linear coordinate change, we may assume that X is defined by the equation

$$x_0^dy_0+x_0^{d-1}x_1y_1+\ldots+x_1^dy_d=0 \tag{2.2}$$

Note that the fibers of the first projection are hyperplanes in $\mathbb{P}^n$, so X is a projective bundle over $\mathbb{P}^1$, a so-called rational normal scroll. In fact, we will show that when X is general, it is isomorphic to $\mathbb{P}(\mathscr{E})$ where $\mathscr{E}=\mathscr{O}_{\mathbb{P}^1}(1)^d\oplus\mathscr{O}_{\mathbb{P}^1}^{n-d}$. Indeed, there is an exact sequence of bundles on $\mathbb{P}^1$

$$0\to\mathscr{O}_{\mathbb{P}^1}(-d)\to\mathscr{O}_{\mathbb{P}^1}^{n+1}\to\mathscr{E}\to 0,$$

which gives an embedding $X=\mathbb{P}(\mathscr{E})\to\mathbb{P}(\mathscr{O}^{n+1})=\mathbb{P}^1\times\mathbb{P}^n$ as a hypersurface. More explicitly, if we define $\phi:\mathbb{P}(\mathscr{E})\to\mathbb{P}^1\times\mathbb{P}^n$ by

$$\phi^*(y_i)=\begin{cases}-x_1z_1 & \text{for } i=0\\ x_0z_i-x_1z_{i+1}, & \text{for } 0<i<d\\ x_0z_d, & i=d\\ z_i & \text{for } i=d+1\ldots n,\end{cases} \tag{2.3}$$

then the image in $\mathbb{P}^1\times\mathbb{P}^n$ is precisely the hypersurface defined by the equation (2.2). Here we consider a basis $z_1,\ldots,z_d$ is a basis for $H^0(\mathbb{P}^1,\mathscr{E}(-1))\subset H^0(\mathbb{P}(\mathscr{E}),\mathscr{O}(1)\otimes\pi^*\mathscr{O}(-1))$ and $z_{d+1},\ldots,z_n$ a basis for $H^0(\mathbb{P}^1,\mathscr{O}^{n-d})\subset H^0(\mathbb{P}(\mathscr{E}),\mathscr{O}(1))$. Note that in terms of the basis H_1,H_2 we have $H_1\sim\pi^*\mathscr{O}_{\mathbb{P}^1}(1)$ and $H_2\sim\mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)$.

The vector bundle $\mathscr{E}$ is generated by the $n+d$ sections $u_{ij}=x_iz_j$ for $i=0,1,j=1,\ldots,d$ and $u_1=z_{d+1}\ldots,u_{n-d}=z_n$. These sections give an embedding of X inside $\mathbb{P}^1\times\mathbb{P}^{n+d-1}$ and the defining equations are the minors of the matrix

$$\begin{pmatrix}x_0 & u_{01} & \ldots & u_{0d}\\ x_1 & u_{11} & \ldots & u_{1d}\end{pmatrix}$$

From this we see that the second projection $p=\mathrm{pr}_2$ is birational and the image Z is a cone over the Segre embedding of $\mathbb{P}^1\times\mathbb{P}^{d-1}$ inside $\mathbb{P}^{n+d-1}$. In fact, from the defining equations it is clear that X is the blow-up of Z along the ideal (u_{01},u_{11}).

2.1. The case $d=1$.

In this case, the second projection contracts the divisor $z_1=0$ and realizes X as a blow-up of $\mathbb{P}^n$ along the codimension 2 linear space $u_{01}=u_{11}=0$. The exceptional divisor is linearly equivalent to $\mathrm{div}z_1=H_2-H_1$. From this description, it is clear that

$$\mathrm{Nef}(X)=\mathrm{Mov}(X)=\mathbb{R}_{\geq 0}H_1+\mathbb{R}_{\geq 0}H_2 \text{ and } \mathrm{Eff}(X)=\mathbb{R}_{\geq 0}H_1+\mathbb{R}_{\geq 0}(H_2-H_1), \tag{2.4}$$

proving Theorem 1.1 in the case $d=e=1$.

2.2. The case $2\leq d<n$.

Here the ideal (u_{01},u_{11}) has codimension 1 in Z an so $p=\mathrm{pr}_2:X\to Z$ is a small contraction. The variety Z has two rulings (corresponing to each factor in the $\mathbb{P}^1 \times \mathbb{P}^{d-1}$). We can blow up the cone Z along the $\mathbb{P}^{d-1}$ ruling, to get a variety $p^+ : X^+ \to Z$ which embeds in $\mathbb{P}^{d-1} \times \mathbb{P}^{n+d-1}$ with determinantal equations as above. The second projection $\pi^+ = \mathrm{pr}_2^+ : X^+ \to \mathbb{P}^{d-1}$ defines another projective bundle, namely $X^+ = \mathbb{P}(\mathscr{O}_{\mathbb{P}^{d-1}}(1)^2 \oplus \mathscr{O}_{\mathbb{P}^{d-1}}^{n-d}) \to \mathbb{P}^{d-1}$ and the induced birational map $X \dashrightarrow X^+$ is the $(-H_1)$-flip of the morphism $p : Y \to Z$. In particular, we see that the transform of $H_2 - H_1$ is nef on X^+. Hence the two extremal contractions of X^+ are given by $X^+ \to Z$ and the bundle projection $\pi^+ : X^+ \to \mathbb{P}^{d-1}$.

Remark 2.1. For $d = 2$ and $n = 3$, this gives the classical Atiyah flop, where the variety Z above is the quadric cone in $\mathbb{P}^4$ and $X \to Z$ and $X^+ \to Z$ are its two small resolutions.

In all, we see that the nef cone of X is spanned by H_1 and H_2, since these are nef, but not ample, and $\mathrm{Eff}(X)$ is spanned by H_2 and $H_2 - H_1$, since these are effective but not big. Moreover, the last two line bundles are movable, and their transforms are nef on X^+. Hence there is a decomposition

$$\mathrm{Eff}(X) = \mathrm{Mov}(X) = \mathrm{Mov}(X) = \mathrm{Nef}(X) \cup \mathrm{Nef}(X^+) = \mathbb{R}_{\geq 0} H_1 + \mathbb{R}_{\geq 0}(H_2 - H_1) \tag{2.5}$$

This is shown in Figure 1.



FIGURE 1. The decomposition of the movable cone

2.3. **The case $d = n$.** In this case, we claim that $Y \simeq \mathbb{P}^1 \times \mathbb{P}^{n-1}$. Indeed, if x_0, x_1 are coordinates on the $\mathbb{P}^1$ and $z_0, \dots, z_{n-1}$ coordinates on the $\mathbb{P}^{n-1}$, the morphism ϕ defined in (2.3) gives an embedding $Y \to \mathbb{P}^1 \times \mathbb{P}^n$ as a bidegree $(n, 1)$ hypersurface. Since $Y \simeq \mathbb{P}^1 \times \mathbb{P}^{n-1}$ and $\operatorname{div} z_i = H_2 - H_1$, we have

$$\mathrm{Eff}(X) = \mathrm{Mov}(X) = \mathrm{Nef}(X) = \mathbb{R}_{\geq 0} H_1 + \mathbb{R}_{\geq 0}(H_2 - H_1) \tag{2.6}$$

and the two contractions are given by the two projections to each factor.

Remark 2.2. Inverting the expressions (2.3) for the z_i, we can compute an explicit basis for $H^0(X, H_2 - H_1)$ in terms of the coordinates on $\mathbb{P}^1 \times \mathbb{P}^n$. Abusing notation, we write

$$\begin{aligned} z_1 &= -y_0/x_1 \\ z_2 &= -\left(x_0 y_0 + x_1 y_1\right)/x_1^2 \\ &\vdots \\ z_d &= -\left(x_0^{d-1} y_0 + \ldots + x_1^{d-1} y_{d-1}\right)/x_1^d = y_d/x_0. \end{aligned}$$

Note for example that z_1 is a global section, since it can only have a pole along the divisor x_1 and modulo the form (2.2), we have $-y_0/x_1 = \left(x_0^{n-1} y_0 + \ldots + x_1^{n-1} y_{n-1}\right)/x_0^n$.

2.4. **Higher degree hypersurfaces.** Let $f_0, \dots, f_d$ be general degree e forms in $y_0, \dots, y_n$ and consider a hypersurface X defined by an equation of the form

$$f = x_0^d f_0 + \dots + x_1^d f_d = 0$$

Let Y be the hypersurface of bidegree $(d, 1)$ defined by the form (2.2) in $\mathbb{P}^1 \times \mathbb{P}^n$. Choose general degree e-forms $f_{d+1}, \dots, f_n$ and define $\eta : \mathbb{P}^n \to \mathbb{P}^n$ to be the morphism defined by $\eta^*(y_i) = f_i$ for $i = 0, \dots, n$. The point of this is that $X = \phi^{-1}(Y)$ where $\phi = (\mathrm{id} \times \eta) : \mathbb{P}^1 \times \mathbb{P}^n \to \mathbb{P}^1 \times \mathbb{P}^n$. Since ϕ is finite and surjective, we also get

$$\mathrm{Eff}(X) = \phi^* \mathrm{Eff}(Y), \quad \mathrm{Mov}(X) = \phi^* \mathrm{Mov}(Y) \text{ and } \mathrm{Nef}(X) = \phi^* \mathrm{Nef}(Y). \tag{2.7}$$

Write for simplicity, $h_i = \mathrm{pr}_i^* \mathscr{O}(1)|_Y$ on Y and $H_i = \mathrm{pr}_i^* \mathscr{O}(1)|_X$ on X. By construction, we have $\phi^* h_1 = H_1$ and $\phi^* h_2 = eH_2$.

Proof of Theorem 1.1(i). By (2.4) and (2.7), we have $\mathrm{Nef}(X) = \mathrm{Mov}(X) = \mathbb{R}_{\geq 0} H_1 + \mathbb{R}_{\geq 0} H_2$ and $\mathrm{Eff}(X) = \mathbb{R}_{\geq 0} H_1 + \mathbb{R}_{\geq 0}(-H_1 + eH_2)$. Moreover, X is the blow-up of $\mathbb{P}^n$ along the complete intersection subvariety $\{f_0 = f_1 = 0\}$, and $eH_2 - H_1$ is linearly equivalent to the exceptional divisor. □

Proof of Theorem 1.1(ii). By (2.5) and (2.7), we have $\mathrm{Nef}(X) = \mathbb{R}_{\geq 0} H_1 + \mathbb{R}_{\geq 0} H_2$ and $\mathrm{Eff}(X) = \mathrm{Mov}(X) = \mathbb{R}_{\geq 0} H_1 + \mathbb{R}_{\geq 0}(-H_1 + eH_2)$. It remains to describe the decomposition of $\mathrm{Mov}(X)$.

Let $p : Y \to Z$ be the small contraction determined by h_2, and let $p^+ : Y^+ \to Z$ be the $(-h_1)$-flip of p. We get a small contraction $q : X \to W$ by taking the Stein factorization of $\eta \circ \phi : X \to Y \to Z$. Since f is finite and surjective, the result [12, Corollary 6.9] says that there exists a $(-H_1)$-flip $X^+ \to W$ of q. Moreover, since $h_2 - h_1$ defines a contraction of Y^+ to a $(d-1)$-dimensional variety, so does $\phi^*(h_2 - h_1) = eH_2 - H_1$ on X^+. Hence the nef cone of X^+ is spanned by H_2 and $eH_2 - H_1$ and we find that $\mathrm{Mov}(X) = \mathrm{Nef}(X) \cup \mathrm{Nef}(X^+) = \mathbb{R}_{\geq 0} H_1 + \mathbb{R}_{\geq 0}(eH_2 - H_1)$. □

Proof of Theorem 1.1(iii). From (2.6) and (2.7), we get $\mathrm{Eff}(X) = \mathrm{Mov}(X) = \mathrm{Nef}(X) = \mathbb{R}_{\geq 0} H_1 + \mathbb{R}_{\geq 0}(-H_1 + eH_2)$ verifying the first part of the theorem. Moreover, the divisor $eH_2 - H_1 = \phi^*(h_2 - h_1)$ is base-point free and determines a morphism to $\mathbb{P}^{n-1}$ given by the composition $X \to Y \to \mathbb{P}^{n-1}$. □

Remark 2.3. For hypersurfaces X of bidegree $(2, e)$ in $\mathbb{P}^1 \times \mathbb{P}^n$, it turns out that the birational model X^+ is actually isomorphic to X. We explain this fact as follows.

Suppose that X is defined by $f(x_0, x_1, y) = x_0^2 f_0(y) + x_0 x_1 f_1(y) + x_1^2 f_2(y) = 0$ in $\mathbb{P}^1 \times \mathbb{P}^n$. Note that the second projection $\mathrm{pr}_2 : X \to \mathbb{P}^n$ is generically 2:1, but it contracts the codimension 2 locus given by $W = \{f_0 = f_1 = f_2 = 0\}$ which is a union of rational curves. Let $\phi : X \to Z$ be the Stein factorization of pr_2. Here Z is the double cover of $\mathbb{P}^n$ branched over the divisor given by $\{f_1^2 - 4 f_0 f_2 = 0\}$ in $\mathbb{P}^n$. Let $\sigma : Z \to Z$ be the natural involution that interchanges the sheets of the double cover. σ induces a birational pseudoautomorphism of X defined outside W. Explicitly, σ sends a point $(x_0, x_1, y) \in X$, to $(x_0', x_1', y) \in X$ where (x_0', x_1', y) is the 'other solution' to the equation $f(x_0, x_1, y) = 0$. Using this description, it is easy to show that $\sigma^* H_1 + H_1 = eH_2$, σ is the $-H_1$-flip of ϕ. This recovers the decomposition of the movable cone $\mathrm{Mov}(X) = \mathrm{Nef}(X) \cup \sigma^* \mathrm{Nef}(X) = \mathbb{R}_{\geq 0} H_1 + \mathbb{R}_{\geq 0}(eH_2 - H_1)$.

3. A GEOMETRIC DESCRIPTION OF THE EXTREMAL CONTRACTIONS

Consider a hypersurface X in $\mathbb{P}^1 \times \mathbb{P}^n$ of bidegree (d, e) with $2 \leq d \leq n$. From the previous section we know that X is a Mori dream space and that the movable cone decomposes as $\mathrm{Nef}(X) \cup \mathrm{Nef}(X^+)$ for some birational model X^+. In this section we give a geometric construction of X^+ and the birational map $X \dashrightarrow X^+$. The idea is to construct an embedding of the hypersurface X inside a toric variety Y so that the birational map $X \dashrightarrow X'$ is induced by a birational map of Y. This will also allow us to give an explicit presentation of the Cox ring of X as a quotient of the Cox ring of Y.

Consider the vector bundle $\mathscr{E} = \mathscr{O}_{\mathbb{P}^1\times\mathbb{P}^n}(0,e) \oplus \mathscr{O}_{\mathbb{P}^1\times\mathbb{P}^n}(1,0)^d$ on $\mathbb{P}^1 \times \mathbb{P}^n$ and its associated variety of hyperplanes $\pi : \mathbb{P}(\mathscr{E}) \to \mathbb{P}^1 \times \mathbb{P}^n$. Consider also the vector bundle map $\mathscr{O}^{d+1}_{\mathbb{P}^1\times\mathbb{P}^n} \to \mathscr{E}$ given by the matrix

$$M = \begin{pmatrix} f_0 & f_1 & \cdots & f_{d-1} & f_d \\ -x_1 & x_0 & 0 & \cdots & 0 \\ 0 & -x_1 & x_0 & 0 & \vdots \\ 0 & \ddots & & \ddots & 0 \\ 0 & \cdots & 0 & -x_1 & x_0 \end{pmatrix} \tag{3.1}$$

Note that $\det M = f$ and so X is precisely the degeneracy locus of the map above. From this we see that $L = \operatorname{coker} M|_X$ defines a line bundle on X and the surjection $\mathscr{E}|_X \to L \to 0$ gives rise to an embedding $X \subset \mathbb{P}(\mathscr{E})$.

Let us study the birational structure of $\mathbb{P}(\mathscr{E})$ more closely. The nef cone of $\mathbb{P}(\mathscr{E})$ is spanned by the three line bundles $L_1 = \pi^*\mathscr{O}(1,0), L_2 = \pi^*\mathscr{O}(0,1)$, and $L_3 = \mathscr{O}(1)$. From this we see that there are three extremal contractions of $\mathbb{P}(\mathscr{E})$: $\pi_1 : \mathbb{P}(\mathscr{E}) \to \mathbb{P}^1$; $\pi_2 : \mathbb{P}(\mathscr{E}) \to \mathbb{P}^n$; and $\mu : \mathbb{P}(\mathscr{E}) \to Y$, where μ is birational.

We make this more explicit as follows. The vector bundle $\mathscr{E}$ is generated by the sections

$$u_0 = y_0^e z_0, u_1 = y_0^{e-1} y_1 z_0, \dots, u_N = y_n^e z_0, \quad u_{0i} = x_0 z_i, u_{1i} = x_1 z_i, \quad i = 1, \dots, d$$

where $z_0 \in H^0(\mathscr{E}(0,-e))$ and $z_1, \dots, z_d \in H^0(\mathscr{E}(-1,0))$. These sections determine a surjection $\mathscr{O}^{N+2d+1} \to \mathscr{E} \to 0$ and hence an embedding $\mathbb{P}(\mathscr{E}) \subset \mathbb{P}^1 \times \mathbb{P}^n \times \mathbb{P}^{N+2d}$ where $N = \binom{n+e}{e} - 1$. The three aforementioned contractions are then $\pi_1 = \mathrm{pr}_1, \pi_2 = \mathrm{pr}_2$, and $\mu = (\mathrm{pr}_1 \times \mathrm{pr}_3)$. Note that the third projection pr_3 defines a morphism $\mathbb{P}(\mathscr{E}) \to Z = \operatorname{Proj} H^0(\mathbb{P}^1 \times \mathbb{P}^n, \operatorname{Sym}\mathscr{E})$ which contracts the divisor $z_0 = 0$.

In terms of the sections u_i, u_{jk}, the variety Z has the following interpretation: it is embedded in $\mathbb{P}^{N+2d}$ as the 'join' of two varieties; the Segre embedding $S_{1,d-1} \subset \mathbb{P}^{2d-1}$ of $\mathbb{P}^1 \times \mathbb{P}^{d-1}$ and the e-th Veronese embedding $V_e \subset \mathbb{P}^N$ of $\mathbb{P}^n$. In other words, if $S_{1,d-1}$ is embedded in $\mathbb{P}^{N+2d}$ in the linear space $\mathbb{P}^{2d-1} = \{u_0 = \cdots = u_N = 0\}$ and V_e in $\mathbb{P}^N = \{u_{01} = \cdots = u_{1d} = 0\}$, then as a set, Z consists of all the lines joining a point $S_{1,d-1}$ with a point in V_e. In terms of equations, Z is defined by the quadratic equations defining the Veronese in $u_0, \dots, u_N$ and the binomial quadrics defining the Segre embedding in $u_{01}, \dots, u_{1d}$. In this notation, $\mathbb{P}(\mathscr{E})$ is the blow-up of Z along the ideal $(u_0, \dots, u_N, u_{01}, u_{11})$ and Y is the blow-up along (u_{01}, u_{11}). The morphism $\nu : Y \to Z$ is small since (u_{01}, u_{11}) has codimension one.

On the other hand, the variety Z can also be blown up along the ideal $(u_{01}, \dots, u_{0d})$ to get a different small resolution $Y^+ \to Z$. If we blow up $(u_0, \dots, u_N, u_{01}, \dots, u_{0d})$, we can identify the blow-up with a projective bundle $\mathbb{P}(\mathscr{E}^+)$ where $\mathscr{E}^+$ is the vector bundle $\mathscr{O}_{\mathbb{P}^{d-1}\times\mathbb{P}^n}(0,e) \oplus \mathscr{O}_{\mathbb{P}^{d-1}\times\mathbb{P}^n}(1,0)^2$ over $\mathbb{P}^{d-1} \times \mathbb{P}^n$. As above, we get an embedding $\mathbb{P}(\mathscr{E}^+) \subset \mathbb{P}^{d-1} \times \mathbb{P}^n \times \mathbb{P}^{N+2d}$ and a contraction $\mathbb{P}(\mathscr{E}^+) \to Y^+$. These varieties fit into the following diagram

$$\begin{tikzcd} \mathbb{P}(\mathscr{E}) \arrow[rr, dashed, "\tilde{\phi}"] \arrow[d, "\mu"'] & & \mathbb{P}(\mathscr{E}^+) \arrow[d, "\mu^+"] & \\ Y \arrow[rr, dashed, "\phi"] \arrow[dr, "\nu"'] & & Y^+ \arrow[dl, "\nu^+"] \arrow[dr, "\mathrm{pr}_1^+"] & \\ & Z & & \mathbb{P}^{d-1} \end{tikzcd} \tag{3.2}$$

where all the morphisms except pr_1^+ are birational.

The variety Y is a $\mathbb{Q}$-factorial toric variety with Picard number 2. Concretely, it is defined in $\mathbb{P}^1 \times \mathbb{P}^{N+2d}$ by two sets of equations: (1): The Veronese quadrics in $u_0, \dots, u_N$; (2) the minors of the $2 \times (d+1)$ matrix

$$\begin{pmatrix} x_0 & u_{01} & \cdots & u_{0d} \\ x_1 & u_{11} & \cdots & u_{1d} \end{pmatrix}$$

We choose a basis $\mathbf{e}_1 = \mu_* L_1$ and $\mathbf{e}_2 = \mu_* L_2$ for $N^1(Y)$. The Cox ring of Y is given by

$$\mathcal{R}(Y) = k[x_0, x_1, y_0, \dots, y_n, z_1, \dots, z_d]$$

where the variables are graded by $\deg x_i = \mathbf{e}_1, \deg y_j = \mathbf{e}_2, \deg z_i = e\mathbf{e}_2 - \mathbf{e}_1$. The variety Y^+ is also toric and has a similar determinantal representation in $\mathbb{P}^{d-1} \times \mathbb{P}^{N+2d}$. Moreover, since ϕ is an isomorphism in codimension 1, there is a graded isomorphism $\mathcal{R}(Y^+) \simeq \mathcal{R}(Y)$.

We know the nef cone of $\mathbb{P}(\mathscr{E})$ and that $\mathbb{P}(\mathscr{E}) \to Y$ contracts the divisor $\{z_0 = 0\}$, so we find that $\operatorname{Nef}(Y) = \mathbb{R}_{\geq 0}\mathbf{e}_1 + \mathbb{R}_{\geq 0}\mathbf{e}_2$. Moreover, $\mathbf{e}_2$ defines the map $\nu : Y \to Z$ and $\mathbf{e}_1$ is ν-ample. If we use the strict transforms of $\mathbf{e}_1, \mathbf{e}_2$ as bases of $N^1(Y^+)$, we have $\operatorname{Nef}(Y^+) = \mathbb{R}_{\geq 0}\mathbf{e}_2 + \mathbb{R}_{\geq 0}(e\mathbf{e}_2 - \mathbf{e}_1)$, and $-\mathbf{e}_1$ is ν^+-ample. Hence $\phi : Y \dashrightarrow Y^+$ is the $(-\mathbf{e}_1)$-flip of ν.

The hypersurface X can be embedded in Y via μ. In fact, from the matrix description (3.1), we see that X is defined in Y by the $d+1$ equations

$$f_0 - x_1 z_1 = 0, f_1 + x_0 z_1 - x_1 z_2 = 0, \dots, f_{d-1} + x_0 z_{d-1} - x_1 z_d = 0, f_d + x_0 z_d = 0. \tag{3.3}$$

The $(-\mathbf{e}_1)$-flip $\phi : Y \dashrightarrow Y^+$ restricts to a rational map $\phi|_X : X \dashrightarrow X^+$. Here X^+ is the image of X under ϕ, which is defined by the equations (3.3) in Y^+. Note that the morphism $\nu|_X$ is small, since $\nu : Y \to Z$ is small and since X is defined as a complete intersection in Y intersecting $\operatorname{Exc}(\nu)$ transversely. Moreover, since $\mathbf{e}_1$ is ν-ample on Y, H_1 is $\nu|_X$-ample on X. Similarly, $-H_1$ is ν^+-ample on X^+. Hence $\phi|_X$ is the $-H_1$-flip of $\nu|_X$. This gives

Proposition 3.1. *The movable cone decomposes as* $\operatorname{Mov}(X) = \operatorname{Nef}(X) \cup \phi|_X^* \operatorname{Nef}(X^+)$.

In fact, with a bit more work we can give a presentation of the Cox ring of X:

Theorem 3.2. *Let X be a general bidegree (d, e) hypersurface in $\mathbb{P}^1 \times \mathbb{P}^n$ with $d \leq n$, defined by the form* (2.1)*. Then*

$$\mathcal{R}(X) = k[x_0, x_1, y_0, \dots, y_n, z_1, \dots, z_d]/I \tag{3.4}$$

where $I = (f_0 - x_1 z_1, f_1 + x_0 z_1 - x_1 z_2, \dots, f_{d-1} + x_0 z_{d-1} - x_1 z_d, f_d + x_0 z_d)$.

Proof. Let R denote the polynomial ring on the right hand side of (3.4). This is the Cox ring of the toric variety Y and I describes the equations of X inside Y. It is clear that for generic $f_0, \dots, f_d$ the ideal I is a complete intersection and that R/I is normal. (For example, this is readily seen to be true in the special case when $f_i = y_i^e$ for $i = 0, \dots, d$, and so it holds for general f_i since forming a regular sequence is a Zariski open condition.)

By [8] the Krull dimension of $\mathcal{R}(X)$ is $n+2$ ($= \dim X + \operatorname{rank} \operatorname{Pic}(X)$). We have $2+n+1+d$ generators of R and $d+1$ relations which form a regular sequence. Hence $\mathcal{R} \subseteq \mathcal{R}(X)$ is a finite extension of $\operatorname{Pic}(X)$-graded normal domains, hence an isomorphism. ☐

Remark 3.3. Using the Koszul complex, it is possible to show directly that the restriction map $H^0(Y, \mathscr{O}_Y(D)) \to H^0(X, \mathscr{O}_X(D))$ is surjective for any effective divisor D with kernel I_D, using the vanishing of higher line bundle cohomology on Y and $\mathbb{P}(\mathscr{E})$ (see also [16]). This gives another proof of Theorem 3.2.

4. Surfaces in $\mathbb{P}^1 \times \mathbb{P}^2$

Let X be a very general surface in $\mathbb{P}^1 \times \mathbb{P}^2$ of bidegree (d, e). Much of the theory from the previous sections can be used to study the birational structure of X, but some care must be taken because the Picard group of X might be strictly larger than $\mathbb{Z}H_1 \oplus \mathbb{Z}H_2$. However, the Noether-Lefschetz theorem of [17] says that when X is very general in its linear system and $\omega_{\mathbb{P}^1\times\mathbb{P}^2} \otimes \mathscr{O}_{\mathbb{P}^1\times\mathbb{P}^2}(X)$ is globally generated, then we have $\mathrm{Pic}(X) = \mathrm{Pic}(\mathbb{P}^1 \times \mathbb{P}^2)$. Note that this is the case if and only if $d \geq 2$ and $e \geq 3$. For the remaining cases, we can proceed by a case-by-case analysis.

$e = 1$. As before, X is a projective bundle over $\mathbb{P}^1$, that is, X is a Hirzebruch surface. In fact, using the fact that maximally balanced splitting type is an open condition on vector bundles, it is not hard to see that $X \simeq \mathbb{P}^1 \times \mathbb{P}^1$ for d even and $X \simeq \mathbb{P}(\mathscr{O} \oplus \mathscr{O}(1)))$ for d odd.

$d = 1$. Here the situation is drastically different than that of hypersurfaces in higher dimension, because hypersurfaces of bidegree $(1, e)$ hypersurfaces are usually *not* Mori dream spaces. In fact, very general hypersurfaces of bidegree $(1, e)$ can be described as the blow-up of $\mathbb{P}^2$ along the e^2 intersection points of two very general degree d curves. This is known to have infinitely many (-1)-curves for $e \geq 3$, so their effective cones of divisors are not rational polyhedral. Note that in these cases the rank of the Picard group is $e^2 + 1$. For $e = 1, 2$, they are Del Pezzo surfaces and hence Mori dream spaces.

It is interesting to note that in this case it is still true that the subalgebra of $\mathcal{R}(X)$ given by $\mathcal{R}(H_1, H_2) = \bigoplus_{(a,b)\in\mathbb{Z}^2} H^0(X, aH_1 + bH_2)$ is finitely generated. Indeed, the H_1, H_2 are base-point free, so the subalgebra $\mathcal{R}(H_1, H_2)_{nef} = \bigoplus_{a,b\geq 0} H^0(X, aH_1 + bH_2)$ is finitely generated by Zariski's theorem [8, Lemma 2.8]. Hence $\mathcal{R}(X, H_1, H_2)$ is finitely generated, since any section s can be written as $s'e^m$ for some $m \geq 0$ where $s' \in \mathcal{R}(H_1, H_2)_{nef}$ and e is a section that defines the exceptional divisor E.

$d = 2$. Very general hypersurfaces X of bidegree $(2, e)$ in $\mathbb{P}^1 \times \mathbb{P}^2$ are Mori dream spaces. Indeed, if $e = 2$, X is a Del Pezzo surface of degree 4, which is a Mori dream space with Picard number 6.

When $e \geq 3$, the Noether-Lefschetz theorem quoted above gives that $\mathrm{Pic}(X) \simeq \mathrm{Pic}(\mathbb{P}^1 \times \mathbb{P}^2)$. As in Section 2, we can define a finite surjective morphism $\phi : X \to Y$ to a hypersurface of bidegree $(2, 1)$ (thus $Y \simeq \mathbb{P}^1 \times \mathbb{P}^1$). This gives that nef cone is spanned by H_1 and $eH_2 - H_1$ and equals the effective cone. Moreover, H_1 and $eH_2 - H_1$ are both base-point free, so by Zariski's theorem again, the ring $\mathcal{R}(X)$ is finitely generated. In fact, $\mathcal{R}(X)$ has a presentation similar to that in (3.4).

When $d \geq 3$ and $e \geq 2$, a very general surface of bidegree (d, e) is not a Mori dream space. We postpone the proof of this claim to the next section.

5. Non-Mori dream space hypersurfaces

In this section we give examples of bidegree (d, e) hypersurfaces in $\mathbb{P}^1 \times \mathbb{P}^n$ which do not have a closed effective cone and hence are not Mori dream spaces. From the the previous sections we know that this can only happen when $d \geq n + 1$ and $e \geq 2$.

We will use the following idea: We let Y be the variety of hyperplanes in a rank n semistable vector bundle over an elliptic curve C and construct a generically finite morphism $Y \to \mathbb{P}^1 \times \mathbb{P}^n$ with image X_0 a divisor of bidegree (d, e). We will see that such maps can be constructed as long as $d \geq n + 1$ and $e \geq 2$. Note that on Y there are plenty of nef divisors which are not semiample, due to the continuous part of the Picard group. In fact, we may pick a basis of divisors L_1, L_2 for $N^1(Y)$ such that one of the extremal rays of the nef cone does not have an effective multiple. We will use this observation to show that a line bundle on $\mathbb{P}^1 \times \mathbb{P}^n$ which restricts to the extremal ray of the effective cone of a general (d, e) hypersurface is not effective on X_0. By semicontinuity, this will imply that the extremal ray of the nef cone of a general divisor is not semiample, since it has no effective multiple. Hence a general bidegree (d, e) hypersurface is not a Mori dream space.

We first need the following lemma which gives a bound for the effective cone of a very general hypersurface in $\mathbb{P}^1 \times \mathbb{P}^n$. It essentially says that the class $neH_2 - dH_1$, which has self-intersection 0, is pseudoeffective.

Lemma 5.1. *Let X be a divisor of bidegree (d, e) in $\mathbb{P}^1 \times \mathbb{P}^n$. Then the* $\operatorname{Eff}(X)$ *contains the subcone*

$$\mathbb{R}_{>0}H_1 + \mathbb{R}_{>0}(neH_2 - dH_1) \tag{5.1}$$

Proof. Let $L = \mathscr{O}_X(aH_1 + bH_2)$ be a line bundle in the span (5.1). We have $L^n = b^{n-1}(bd + aen) > 0$. Furthermore, from the exact sequence $0 \to \mathscr{O}_{\mathbb{P}^1\times\mathbb{P}^n}(a-d, b-e) \to \mathscr{O}_{\mathbb{P}^1\times\mathbb{P}^n}(a, b) \to L \to 0$ and cohomology vanishing of line bundles on $\mathbb{P}^1 \times \mathbb{P}^n$, we see that $H^i(Y, L) = 0$ for $i > 1$. Hence $h^0(mL) \geq \chi(mL) = m^n L^n/n! + \dots$ and so L is big. ☐

The line bundles in this cone correspond exactly to the line bundles L such that $L^n > 0$ and $L^{n-1} \cdot H_1 > 0$. The two divisors H_1 and $neH_2 - dH_1$ will in fact turn out to be extremal in the effective cone, so generically there exists no divisors of negative self-intersection on X.

5.1. **Construction of the special hypersurfaces.** Let C be a smooth elliptic curve. By results of Atiyah [2], C has a semistable rank $r = n$ vector bundle $\mathscr{E}$ of degree $d > n$. Let $Y = \mathbb{P}(\mathscr{E})$ denote the variety of hyperplanes in $\mathscr{E}$ with its projection $\pi : Y \to C$.

An essential point of the construction is defining two morphisms $f : Y \to \mathbb{P}^1$ and $g : Y \to \mathbb{P}^n$, so that the induced map $F = f \times g : Y \to \mathbb{P}^1 \times \mathbb{P}^n$ is generically injective and the image of F is a divisor of bidegree (d, e). Let $\mathfrak{d} = p_1 + \dots + p_e$ be a divisor of degree e on C. We will define f and g using sections of the line bundles $L_1 = \pi^*\mathscr{O}_C(\mathfrak{d})$ and $L_2 = \mathscr{O}(1)$ on $\mathbb{P}(\mathscr{E})$ and arrange $\mathfrak{d}$ to ensure that this condition holds. Note that $L_1 \cdot L_2^{n-1} = e$ and $L_2^n = d$.

Lemma 5.2. *The line bundles L_1 and L_2 are base-point free and L_2 is ample.*

Proof. L_1 is the pullback of a base-point free divisor on C. L_2 is base-point free because $\mathscr{E}$ is generated by sections: Indeed, this is true for any semistable vector bundle of degree $d > r(2g(C) - 1) = n$. When $\mathscr{E}$ is semistable, any effective divisor on $\mathbb{P}(\mathscr{E})$ is nef [13, 1.5.A]. Moreover, L_2 is big (since $h^i(Y, \mathscr{O}(m)) = h^i(C, S^m\mathscr{E}) = 0$ for all $m > 0$ and $i > 0$, by the semistability of $\mathscr{E}$), so it lies in the interior of the nef cone and hence is ample. ☐

A general base-point free pencil of sections of $\mathscr{O}_C(\mathfrak{d})$ defines a degree e morphism $C \to \mathbb{P}^1$. Composing with π, we see that a pencil of sections of L_1 defines a morphism $f : Y \to \mathbb{P}^1$ whose generic fiber consists of e distinct fibers of π. Similarly, note that if $\mathscr{E}$ is semistable of degree $d > n(2g - 1) = n$, we have $h^1(C, \mathscr{E}) = 0$ and so by Riemann-Roch, $h^0(C, \mathscr{E}) = d \geq n + 1$. A choice of $n + 1$ generic sections defines a finite morphism $g : Y \to \mathbb{P}^n$ of degree d.

Lemma 5.3. *For $\mathfrak{d} \in \operatorname{Sym}^e C$ very general, the image X_0 of the morphism $F = f \times g : X \to \mathbb{P}^1 \times \mathbb{P}^n$ is a divisor of bidegree (d, e).*

Proof. First of all, F is finite, since g is finite, and so the image is a (possibly singular) hypersurface in $\mathbb{P}^1 \times \mathbb{P}^n$. We will show that F has degree 1 below.

Given this, it is clear that the image X_0 has bidegree (d, e). Indeed, note that the projections $\operatorname{pr}_1 : Y \to \mathbb{P}^1$ and $\operatorname{pr}_2 : Y \to \mathbb{P}^n$ factor through $f \times g$ and determine the bidegree uniquely: If X_0 has bidegree (a, b), we have $a = X_0 \cdot \operatorname{pr}_2^* \mathscr{O}_{\mathbb{P}^n}(1)^n = L_2^n = d$ and $b = X_0 \cdot \operatorname{pr}_2^* \mathscr{O}_{\mathbb{P}^1}(1) \cdot \operatorname{pr}_2^* \mathscr{O}_{\mathbb{P}^n}(1)^{n-1} = L_1 \cdot L_2^{n-1} = e$.

We now show that F has degree 1. It is clear that the generic fiber of f consists of e fibers of π and g is finite of degree d. Let $y \in Y$ be a general point. To show that F is generically injective it suffices to show that $f^{-1}(f(y))$ avoids $f^{-1}(f(y'))$ for any $y' \in g^{-1}(g(y)) - \{y\}$. Indeed, if this is true, then we have $F^{-1}(F(y)) = f^{-1}(f(y)) \cap g^{-1}(g(y)) = \{y\}$.

Note that if l is a line through the point $g(y) \in \mathbb{P}^n$, then the preimage $g^{-1}(l)$ is generically a section of π (because $L_2^{n-1} \cdot \pi^*\mathscr{O}_C(p) = 1$ for a fiber over $p \in C$). This means that if y is a general point, the points in the fiber $g^{-1}(g(y))$ must lie in different fibers of π. Now the claim follows, since of generic $\mathfrak{d}$ and f, the preimage $f^{-1}(f(y))$ intersects these fibers only at y. Thus F is generically injective and the proof is complete. ☐

Lemma 5.4. *If $\mathfrak{d}$ is very general, the divisor $D = enL_2 - dL_1$ on Y does not have a positive integral multiple which is effective.*

Proof. We need to show that for each $m > 0$,

$$H^0(Y, \mathscr{O}(mD)) = H^0(C, \pi_*\mathscr{O}(mD)) = H^0(C, S^{enm}\mathscr{E} \otimes \mathscr{O}_C(-dm\mathfrak{d})) = 0$$

Note that $S^{emn}\mathscr{E}$ is a semistable vector bundle of rank $r = \binom{emn+n-1}{n-1}$ and degree $edmr$ and so $S^{emn}\mathscr{E} \otimes \mathscr{O}_C(-dm\mathfrak{d})$ is semistable of degree 0. If $\mathfrak{d}$ is very general, then these vector bundles do not have any global sections by the lemma below. ☐

Lemma 5.5. *Let $\mathscr{E}$ be a semistable vector bundle of rank r and degree 0 on a curve of positive genus. Then for a line bundle L defining a general point in $Pic^0(C)$, we have $H^0(\mathscr{E} \otimes L) = 0$.*

Proof. If $\mathscr{E}$ is a line bundle, then the result is clear, since the only effective line bundle of degree 0 is the trivial bundle and $\mathrm{Pic}^0(C)$ has dimension > 0. So we may assume that rank $\mathscr{E} \geq 2$.

The result holds if $\mathscr{E}$ is stable, because in that case so is $\mathscr{E} \otimes L$, and if $\mathscr{E} \otimes L$ has a section, then $\mathscr{O}$ is a subsheaf, contradicting the stability condition.

If $\mathscr{E}$ is strictly semistable, then the Jordan-Hölder filtration says that there is a semistable subbundle $\mathscr{E}' \subset \mathscr{E}$ such that $\mathscr{E}/\mathscr{E}'$ is a stable vector bundle and both $\mathscr{E}$ and $\mathscr{E}/\mathscr{E}'$ have the same slope as $\mathscr{E}$ (that is, 0). From this we get an exact sequence of degree 0 vector bundles

$$0 \to \mathscr{E}' \to \mathscr{E} \to \mathscr{E}/\mathscr{E}' \to 0.$$

Tensoring this with L and taking cohomology, the result follows by induction on the rank. ☐

We are ready to prove the main theorem of this section:

Theorem 5.6. *Let X be a very general hypersurface of $\mathbb{P}^1 \times \mathbb{P}^n$ of bidegree (d, e) with $d \geq n+1$ and $e \geq 2$. Then the effective cone of X is not closed. In particular, X is not a Mori dream space.*

Proof. We show that the effective cone is not closed. To do this it is sufficient by semicontinuity of $\dim H^0$ to exhibit a single divisor X_0 of bidegree (d, e) such that no multiple of the line bundle $L := \mathrm{pr}_1^*\mathscr{O}_{\mathbb{P}^1}(-d) \otimes \mathrm{pr}_2^*\mathscr{O}_{\mathbb{P}^n}(ne)$ restricts to an effective divisor on X_0; then the same conclusion holds for a very general deformation of it. Since this line bundle is pseudoeffective on a very general hypersurface, the result follows.

We will let X_0 be the image of $Y = \mathbb{P}(\mathscr{E})$ under the morphism $F = f \times g$ defined earlier. By construction, the image X_0 is a divisor of bidegree (d, e) such that the line bundle $F^*L|_{X_0} = \mathscr{O}_Y(enL_2 - dL_1)$ has no effective multiple on Y (Lemma 5.4). Note that X_0 is reduced (although it may be singular), hence the natural map $\mathscr{O}_{X_0} \to F_*\mathscr{O}_Y$ is injective, and we have

$$H^0(X_0, L|_{X_0}) \subseteq H^0(X_0, L|_{X_0} \otimes F_*\mathscr{O}_Y) = H^0(Y, F^*L|_{X_0}) = 0$$

The same argument shows that no multiple of L is effective on X_0 and the proof is complete. ☐

Corollary 5.7. *Let X be a very general divisor $X \subset \mathbb{P}^1 \times \mathbb{P}^n$ of bidegree (d, e) with $d \geq n+1$ and $e \geq 2$. Then*

$$\overline{\mathrm{Eff}}(X) = \overline{\mathrm{Mov}}(X) = \mathrm{Nef}(X) = \mathbb{R}_{\geq 0}H_1 + \mathbb{R}_{\geq 0}(neH_2 - dH_1). \tag{5.2}$$

Proof. By Lemma 5.1 we have that the pseudoeffectivecone contains the cone on the right hand side of (5.2), and so two cones coincide by the theorem. Moreover, $neH_2 - dH_1$ is nef on the special hypersurface X_0 used in the proof of the theorem, since it pulls back to a nef divisor on $\mathbb{P}(\mathscr{E})$ via a finite surjective morphism. Moreover, on X_0, the pseudoeffective cone and the nef cone coincide, so by the argument of [14, Lemma 4.1], the same conclusion holds for a very general deformation X of X_0. □

Example 5.8. In dimension 2, $(3, 2)$ is the first bidegree for which a very general divisor is not a Mori dream space. This variety is rational surface which is isomorphic to a blow-up of a Hirzebruch surface in 9 general points, so the Picard number is in fact 11.

Example 5.9. By the Noether-Lefschetz theorem, a very general surface of bidegree $(3, 3)$ in $\mathbb{P}^1 \times \mathbb{P}^2$ has Picard number 2. By the theorem, the line bundle $L = 2H_2 - H_1$ is nef, but is not semiample. In fact, L is *strictly nef*, in the sense that $\deg L|_C > 0$ for every curve C (since $L \cdot C = 0$ implies $C \sim_{\mathbb{Q}} L$, and L is not $\mathbb{Q}$-linearly equivalent to an effective divisor). This gives a simple counterpart of Mumford's example mentioned in the introduction.

Example 5.10. A very general divisor of bidegree $(4, 2)$ in $\mathbb{P}^1 \times \mathbb{P}^3$ can be viewed as a quadric surface bundle over $\mathbb{P}^1$. It is therefore a rational threefold with Picard number 2, which is not a Mori dream space.

6. Hypersurfaces in products of several projective spaces

For hypersurfaces X in products of projective spaces with more than one $\mathbb{P}^1$-factor, the situation becomes more complicated. In particular, we don't expect Mori dream spaces, even for low degree hypersurfaces. The following example, which appears in the work of Kawamata [11], illustrates this already for a Calabi-Yau threefold.

Consider a smooth hypersurface of tridegree $(2, 2, 3)$ in $\mathbb{P} = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^2$ defined by an equation

$$f(x_i, y_i, z_i) = x_0^2 f_0 + x_0 x_1 f_1 + x_1^2 f_2$$

where f_0, f_1, f_2 are forms of tridegree $(0, 2, 3)$. Note that the projection $(\mathrm{pr}_1 \times \mathrm{pr}_3) : X \to \mathbb{P}^1 \times \mathbb{P}^2$ contracts the codimension 2 locus $W = \{f_0 = f_1 = f_2 = 0\}$ which is a union of 54 rational curves. Taking the Stein factorization gives a small contraction $\phi : X \to Z$ where Z is the double cover of $\mathbb{P}^1 \times \mathbb{P}^2$ ramified over the divisor defined by $f_1^2 - 4f_0 f_2 = 0$. Note that Z has a natural involution $\sigma' : Z \to Z$, which switches the sheets of the covering. This determines a birational pseudoautomorphism $\sigma : X \dashrightarrow X$ defined outside W. In terms of H_1, H_2, H_3 it is not hard to show that $\sigma^* H_1 + H_1 = 2H_2 + 3H_3$ and that $\sigma : X \dashrightarrow X$ is the $(-H_1)$-flop of ϕ.

One can repeat the argument with the other contraction $(\mathrm{pr}_2 \times \mathrm{pr}_3) : X \to \mathbb{P}^1 \times \mathbb{P}^2$ to get another pseudoautomorphism σ' of X. Moreover, σ and σ' generate an infinite subgroup of the group of pseudoautomorphisms of X, $\mathrm{PsAut}(X)$. In fact, also the group $\mathrm{PsAut}(X)^* = \mathrm{im}(\mathrm{PsAut}(X) \to GL(N^1(X))$ is infinite. In particular, X is not a Mori dream space, because there are infinitely many non-isomorphic marked small $\mathbb{Q}$-factorial modifications.

Using essentially the same technique, one can show the following result:

Proposition 6.1. *Let X be a smooth Calabi-Yau hypersurface of dimension ≥ 3 in $\mathbb{P} = (\mathbb{P}^1)^m \times \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$ where $n_1, \ldots, n_k \geq 2$ and let $H_i = \mathrm{pr}_i^* \mathscr{O}(1)$. Then the nef cone is given by* $\mathrm{Nef}(X) = \mathbb{R}_{\geq 0} H_1 + \mathbb{R}_{\geq 0} H_2 + \ldots + \mathbb{R}_{\geq 0} H_{k+m}$. *Moreover, the following hold:*

- *(i) If $m = 0$, then* $\mathrm{Eff}(X) = \mathrm{Mov}(X) = \mathrm{Nef}(X) = \mathrm{Eff}(\mathbb{P})$ *and X is a Mori dream space.*
- *(ii) If $m = 1$, then the effective cone is strictly larger than that of $\mathbb{P}$, X is a Mori dream space.*
- *(iii) If $m > 1$, then X is not a Mori dream space. In fact the group $PsAut(X)^*$ is infinite and the movable cone is not rational polyhedral.*

Proof. The description of the nef cone follows from a more general theorem of Kollár [3], which states that if Y is a smooth Fano variety of dimension at least 4 and $X \in |-K_Y|$ is a smooth divisor, then the inclusion induces an isomorphism of the cones of curves $i_* : \overline{NE}_1(X) \to \overline{NE}_1(Y)$. Taking duals gives that the nef cone is the first quadrant $\sum_j \mathbb{R}_{\geq 0} H_j$ in $N^1(X)$.

The case (i) follows from the results of Jow and Artebani–Laface, as in Proposition 1.3.

For (ii), let $\sigma : X \dashrightarrow X$ be the pseudoautomorphism of X obtained by viewing X as a double cover of $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$. It is not hard to show that any divisor D lying in the'coordinate planes' $\mathbf{e}_i^\perp = \sum_{j \neq i} \mathbb{R}_{\geq 0} H_j$ is not big for any $i > 1$. This follows for example by taking cohomology of the ideal sheaf sequence $0 \to \mathscr{O}_{\mathbb{P}}(D - X) \to \mathscr{O}_{\mathbb{P}}(D) \to \mathscr{O}_X(D) \to 0$ and using the vanishing of higher cohomology of line bundles on $\mathbb{P}$. Hence the only supporting hyperplane of $\mathrm{Nef}(X)$ containing big divisors is $e_1^\perp$. Note that this hyperplane is fixed under the involution σ^*, and so by applying σ^* to $\mathrm{Nef}(X)$, we see that any divisor in the boundary of $\mathrm{Nef}(X) \cup \sigma^* \mathrm{Nef}(X)$ is not big. Hence $\mathrm{Eff}(X) = \mathrm{Mov}(X) = \mathrm{Nef}(X) \cup \sigma^* \mathrm{Nef}(X)$ and so X is a Mori dream space.

(iii) follows as in the example above, noting that two $\mathbb{P}^1$-factors give rise to two pseudoautomorphisms σ, σ' generating an infinite subgroup of $\mathrm{PsAut}(X)^*$. □

In [5], Cantat and Oguiso give a very detailed overview of the cones of effective, movable, and nef divisors on hypersurfaces in $(\mathbb{P}^1)^m$ of multidegree $(2, \ldots, 2)$. In particular, they verify the Morrison-Kawamata cone conjecture for these hypersurfaces, which means that even though the movable cone itself is not rational polyhedral, it has a rational polyhedral fundamental domain under the action of $\mathrm{PsAut}(X)$ on $N^1(X)$. In fact, they show that $\mathrm{PsAut}(X) \simeq \mathbb{Z}/2\mathbb{Z} * \cdots * \mathbb{Z}/2\mathbb{Z}$, generated by the birational involutions above and that the fundamental domain in question is the nef cone of X. It is likely that these results generalize to the hypersurfaces in Proposition 6.1.

REFERENCES

[1] M. Artebani, A. Laface, Hypersurfaces in Mori dream spaces. J. Algebra 371 (2012), 26–37.

[2] M. F. Atiyah, Vector bundles over an elliptic curve. Proc. London Math. Soc. 7 (1957) 414–452.

[3] C. Borcea, Homogeneous vector bundles and families of Calabi-Yau threefolds. II, Several complex variables and complex geometry, 1989, Proc. Sympos. Pure Math. 52, (1991) 83–91.

[4] C. Birkar and P. Cascini, C. Hacon, J. McKernan, Existence of minimal models for varieties of log general type, J. Amer. Math. Soc., 23, 2010. 405–468.

[5] S. Cantat, K. Oguiso. Birational automorphism groups and the movable cone theorem for Calabi-Yau manifolds of Wehler type via universal Coxeter groups. arXiv:1107.5862

[6] Jürgen Hausen, Cox rings and combinatorics. II, Mosc. Math. J. 8 (2008), no. 4, 711 757, 847

[7] B. Hassett, H. W. Lin, C.-L. Wang, The weak Lefschetz principle is false for ample cones. Asian J. Math. 6, (2002) 95–100.

[8] Y. Hu, Y, S. Keel, Mori dream spaces and GIT. Michigan Math. J. 48 (2000), 331–348.

[9] A. Ito. Examples of Mori dream spaces with Picard number two. arXiv:1304.8005

[10] S.-Y. Jow, A Lefschetz hyperplane theorem for Mori dream spaces. Math. Z. 268 (2011), 197–209.

[11] Y. Kawamata, On the cone of divisors of Calabi-Yau fiber spaces, Internat. J. Math. 8 (1997) 665687.

[12] J. Kollár, S. Mori S., Birational geometry of algebraic varieties, Cambridge tracts in mathematics, vol. 134, Cambridge University Press, 1998.

[13] R. Lazarsfeld, *Positivity in algebraic geometry I, II*, in Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. 48-49. Springer-Verlag, Berlin, 2004.

[14] C. Mourougane, Families of hypersurfaces of large degree. J. Eur. Math. Soc.14 (2012), 911–936.

[15] M. Nagata, On rational surfaces. II. Mem. Coll. Sci. Univ. Kyoto Ser. A Math. 33 (1960/1961), 271–293.

[16] J. C. Ottem, The Cox ring of a K3 surface of Picard number two. J. of Pure and Applied Algebra. 217 (2013), 709–715

[17] G. V. Ravindra, V. Srinivas, The Noether-Lefschetz theorem for the divisor class group. J. Algebra 322 (2009), no. 9, 3373–3391.

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, UNIVERSITY OF CAMBRIDGE, WILBERFORCE ROAD, CAMBRIDGE CB3 0WB, UK

E-mail address: `J.C.Ottem@dpmms.cam.ac.uk`